



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Introducing an algal carbon-concentrating mechanism into higher plants: location and incorporation of key components

Citation for published version:

Atkinson, N, Feike, D, Mackinder, LCM, Meyer, MT, Griffiths, H, Jonikas, MC, Smith, AM & McCormick, AJ 2015, 'Introducing an algal carbon-concentrating mechanism into higher plants: location and incorporation of key components', *Plant biotechnology journal*. <https://doi.org/10.1111/pbi.12497>

Digital Object Identifier (DOI):

[10.1111/pbi.12497](https://doi.org/10.1111/pbi.12497)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Plant biotechnology journal

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Introducing an algal carbon-concentrating mechanism into higher plants: location and incorporation of key components

Nicky Atkinson¹, Doreen Feike², Luke C. M. Mackinder³, Moritz T. Meyer⁴, Howard Griffiths⁴, Martin C. Jonikas³, Alison M. Smith² and Alistair J. McCormick^{1,2,*}

¹SynthSys & Institute of Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

²Department of Metabolic Biology, John Innes Centre, Norwich Research Park, Norwich, UK

³Department of Plant Biology, Carnegie Institution for Science, Stanford, CA, USA

⁴Department of Plant Sciences, University of Cambridge, Cambridge, UK

Received 1 July 2015;

revised 18 September 2015;

accepted 29 September 2015.

*Correspondence (Tel +44 (0)1316505316;

fax +44 (0) 1316508650;

email alistair.mccormick@ed.ac.uk)

Summary

Many eukaryotic green algae possess biophysical carbon-concentrating mechanisms (CCMs) that enhance photosynthetic efficiency and thus permit high growth rates at low CO₂ concentrations. They are thus an attractive option for improving productivity in higher plants. In this study, the intracellular locations of ten CCM components in the unicellular green alga *Chlamydomonas reinhardtii* were confirmed. When expressed in tobacco, all of these components except chloroplastic carbonic anhydrases CAH3 and CAH6 had the same intracellular locations as in *Chlamydomonas*. CAH6 could be directed to the chloroplast by fusion to an Arabidopsis chloroplast transit peptide. Similarly, the putative inorganic carbon (Ci) transporter LCI1 was directed to the chloroplast from its native location on the plasma membrane. CCP1 and CCP2 proteins, putative Ci transporters previously reported to be in the chloroplast envelope, localized to mitochondria in both *Chlamydomonas* and tobacco, suggesting that the algal CCM model requires expansion to include a role for mitochondria. For the Ci transporters LCIA and HLA3, membrane location and Ci transport capacity were confirmed by heterologous expression and H¹⁴CO₃⁻ uptake assays in *Xenopus* oocytes. Both were expressed in Arabidopsis resulting in growth comparable with that of wild-type plants. We conclude that CCM components from *Chlamydomonas* can be expressed both transiently (in tobacco) and stably (in Arabidopsis) and retargeted to appropriate locations in higher plant cells. As expression of individual Ci transporters did not enhance Arabidopsis growth, stacking of further CCM components will probably be required to achieve a significant increase in photosynthetic efficiency in this species.

Keywords: photosynthesis improvement, carbon-concentrating mechanism, bicarbonate transporter, *Arabidopsis thaliana*, tobacco, *Chlamydomonas reinhardtii*.

Introduction

Most plants, including the major grain crops rice and wheat, assimilate carbon using the C₃ photosynthetic pathway. C₃ plants rely on passive diffusion to deliver carbon dioxide (CO₂) from the atmosphere (ca. 400 ppm) to the chloroplasts inside leaf mesophyll cells, wherein CO₂ photoassimilation proceeds via the primary carboxylase enzyme, ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO, EC 4.1.1.39). Diffusive resistances result in a gradient (ca. 40% under high irradiance) between CO₂ levels in the substomatal cavity of the leaf and the steady-state level of dissolved CO₂ in chloroplasts (10–20 µM at 25 °C) (Price *et al.*, 2011). Here, CO₂ is not saturating for RuBisCO and oxygen (O₂; ca. 250 µM at 25 °C) competes at the RuBisCO active sites, resulting in both loss of assimilated carbon and nitrogen and energy consumption in the photorespiratory pathway that recycles the product of RuBP oxygenation (Sharkey, 1988). The productivity of C₃ crops is thus limited by the efficiency of CO₂ photoassimilation, even when grown under elevated CO₂ levels (up to 650 ppm) (Long *et al.*, 2006). Generating C₃ crop plants

with increased photosynthetic efficiencies is a major target for improving yields and safeguarding future food security. Strategies under consideration and development include modifying canopies to increase light interception, enhancing repair mechanisms to overcome lags associated with photoprotection, increasing the efficiency of RuBisCO and eliminating photorespiration by introducing molecular components of microbial carbon-concentrating mechanisms (CCM) (Lin *et al.*, 2014a,b; Long *et al.*, 2015; Parry *et al.*, 2013; Whitney *et al.*, 2011; Zhu *et al.*, 2010).

Many photosynthetic organisms including cyanobacteria, most green algae and a single group of land plants, the hornworts, have evolved biophysical CCMs that actively increase the CO₂ concentration around RuBisCO, thus suppressing RuBisCO oxygenase activity and associated photorespiration. In eukaryotic algae, CCMs involve inorganic carbon (Ci) transporters at the plasma membrane and chloroplast envelope and carbonic anhydrases, which act in concert to deliver above ambient concentrations of CO₂ to RuBisCO, usually within a chloroplast microcompartment called the pyrenoid. The pyrenoid is mainly composed of densely packaged RuBisCO (Engel *et al.*, 2015).

Please cite this article as: Atkinson, N., Feike, D., Mackinder, L.C.M., Meyer, M.T., Griffiths, H., Jonikas, M.C., Smith, A.M. and McCormick, A.J. (2015) Introducing an algal carbon-concentrating mechanism into higher plants: location and incorporation of key components. *Plant Biotechnol. J.*, doi: 10.1111/pbi.12497

Table 1 Chlamydomonas CCM genes used in this study. Locus name refers to the gene ID as supplied by Phytozome v5.5 (http://phytozome.jgi.doe.gov/pz/portal.html#?info?alias=Org_Creinhardtii)

Gene	Locus name	Protein length, size of precursor	Putative function in native algal CCM	Examples of experimental evidence for function	Mutant phenotype
<i>HLA3</i>	Cre02.g097800	1325 aa, 147 kDa	Ci uptake into cytosol	RNAi lines grown under alkaline conditions (main Ci species is HCO_3^-) have a HCR (1); synergistic effect with <i>LCIA</i> (1;2); ^{14}Ci uptake assay (2;3); affinity for Ci when controlling gene expression with dTALE (3); + work presented here	Reduction in Ci accumulation when $\text{CO}_2 < 0.02\%$; HCR
<i>LCI1</i>	Cre03.g162800	192 aa, 21 kDa	Ci uptake into cytosol	^{14}Ci uptake assay (4)	No published mutant; overexpression in CCM regulatory mutant promotes HCO_3^- uptake
<i>LCIA</i>	Cre06.g309000	336 aa, 35 kDa	Ci transport from cytosol to stroma	Synergistic effect with <i>HLA3</i> (2;3); electrophysiology in <i>Xenopus</i> oocytes (5); reduced affinity for Ci at alkaline pH (6); + work presented here	Reduction in Ci accumulation when $\text{CO}_2 < 0.02\%$; HCR
<i>CCP1</i>	Cre04.g223300	358 aa, 38 kDa	Ci transport from cytosol to stroma	RNAi lines support role in Ci transport (7); putative localization inferred bioinformatically	No published mutant
<i>CCP2</i>	Cre04.g222750	355 aa, 38 kDa	Ci transport from cytosol to stroma	RNAi lines support role in Ci transport (7); putative localization inferred bioinformatically	No published mutant
<i>CAH1</i>	Cre04.g223100	377 aa, 42 kDa	CO_2 and HCO_3^- at cell surface	Absence of growth effect in the presence of membrane impermeable CA inhibitors (9)	No apparent deleterious effect on growth
<i>CAH6</i>	Cre12.g485050	264 aa, 28 kDa	Recapture of CO_2 leaking from the pyrenoid	No experimental evidence of function; putative role in CCM inferred from putative localization	No published mutant
<i>CAH3</i>	Cre09.g415700	310 aa, 33 kDa	Terminal dehydration of HCO_3^- to CO_2 , to saturate RuBisCO in the pyrenoid	Low CO_2 -induced phosphorylation relocates <i>CAH3</i> preferentially to pyrenoid tubules (13); alternative CCM-unrelated function (regulation of water oxydation at PSII) has been proposed (14)	Mutation produces overaccumulation of Ci
<i>LCIB</i>	Cre10.g452800	448 aa, 48 kDa	CO_2 uptake or trapping of stromal CO_2 , pyrenoid localization	Synergetic role with Ci pumps (<i>lciA/lciB</i> double mutant lethal when $\text{CO}_2 < 0.02\%$) (1); MS-identification of <i>LCIB</i> -FLAG pull-down and gel filtration showed that <i>LCIB</i> -C form a 360 kDa hetero-hexamer, which localizes around the pyrenoid when $\text{CO}_2 < 0.02\%$ (16)	Lethal under air-level CO_2 but rescued when $\text{CO}_2 < 0.02\%$
<i>LCIC</i>	Cre06.g307500	443 aa, 48 kDa	CO_2 uptake or trapping of stromal CO_2 , pyrenoid localization	MS-identification of <i>LCIB</i> -FLAG pull-down and gel filtration showed that <i>LCIB</i> -C form a 360 kDa hetero-hexamer, which localizes around the pyrenoid when $\text{CO}_2 < 0.02\%$ (16)	No published mutant
Gene	Locus name	Functional annotation	Location as per latest CCM model (Wang <i>et al.</i> , 2015)	Examples of experimental evidence for location; work presented here for all ten genes	References
<i>HLA3</i>	Cre02.g097800	ABC transporter superfamily	Plasmamembrane	Immunofluorescence (2); immunoblot on membrane fractions (2;3)	(1) Duanmu <i>et al.</i> (2009a); (2) Yamano <i>et al.</i> (2015); (3) Gao <i>et al.</i> (2015)
<i>LCI1</i>	Cre03.g162800	Unknown	Plasmamembrane	Immunofluorescence + GFP fusion + immunoblot on membrane fractions (4)	(4) Ohnishi <i>et al.</i> (2010)
<i>LCIA</i>	Cre06.g309000	Formate/nitrite transporter	Chloroplast membrane	Immunofluorescence (2;6)	(5) Mariscal <i>et al.</i> (2006); (6) Wang and Spalding, (2014a)
<i>CCP1</i>	Cre04.g223300	Mitochondrial carrier protein	Chloroplast membrane		(7) Pollock <i>et al.</i> (2004); (8) Ramazanov <i>et al.</i> (1993)

Table 1 Continued

Gene	Locus name	Functional annotation	Location as per latest CCM model (Wang <i>et al.</i> , 2015)	Examples of experimental evidence for location; work presented here for all ten genes	References
				Immunoblot on membrane fractions (8); + work presented here	
CCP2	Cre04.g222750	Mitochondrial carrier protein	Chloroplast membrane	Immunoblot on membrane fractions (8); + work presented here	(7) Pollock <i>et al.</i> (2004); (8) Ramazanov <i>et al.</i> (1993)
CAH1	Cre04.g223100	α carbonic anhydrase	Periplasmic space	Cell wall lysis + CA assay on supernatant (10); immunogold (11)	(9) Moroney and Tolbert, (1985); (10) Kimpel <i>et al.</i> (1983); (11) Ynalvez <i>et al.</i> (2008)
CAH6	Cre12.g485050	β carbonic anhydrase	Stroma	Immunogold (12)	(12) Mitra <i>et al.</i> (2004)
CAH3	Cre09.g415700	α carbonic anhydrase	Thylakoid lumen	Immunogold (12;15)	(13) Blanco-Rivero <i>et al.</i> (2012); (14) Villarejo <i>et al.</i> (2002); (15) Markelova <i>et al.</i> (2009)
LCIB	Cre10.g452800	Unknown	Stroma	Immunogold (16); GFP fusion (16,17); immunofluorescence (6;18)	(16) Yamano <i>et al.</i> (2010); (17) Yamano <i>et al.</i> (2014); (18) Wang and Spalding (2014b)
LCIC	Cre06.g307500	Unknown	Stroma	Immunogold + immunofluorescence (16)	

CA, carbonic anhydrase; Ci, inorganic carbon; HCR: high CO₂-requiring phenotype, a nonlethal mutation rescued by growth under elevated CO₂ (2–5% [v/v]).

Whilst not all algae with a CCM have a pyrenoid, the microcompartment enhances the efficiency of CO₂ assimilation (Morita *et al.*, 1998). Theoretical modelling approaches have demonstrated the requirement for a pyrenoid in algal systems (Badger *et al.*, 1998) and shown that some form of microcompartment containing RuBisCO would also be needed for a successful CCM in higher plant systems (Price *et al.*, 2013).

The best-characterized algal CCM is that of the model green alga *Chlamydomonas reinhardtii* (*Chlamydomonas* throughout). To date, a large number of molecular components have been implicated in the *Chlamydomonas* CCM through mutant screens, transcriptomic studies and functional homology with components in other photosynthetic organisms (Brueggeman *et al.*, 2012; Fang *et al.*, 2012; Meyer and Griffiths, 2013). At least 14 genes are thought to be important in maintaining a fully functional CCM under ambient or below ambient CO₂ concentrations (see Wang *et al.* (2015) for review). Briefly, these include five Ci transporters, four CAs, two pyrenoid peripheral proteins, a putative methyl transferase and two nuclear transcription regulators. The precise function and importance of these components remains only partly understood. In addition, the *Chlamydomonas* CCM requires a RuBisCO that can be targeted to the pyrenoid, a property known to be dependent on sequence elements of the RuBisCO small subunit (Genkov *et al.*, 2010; Meyer *et al.*, 2012).

To examine the feasibility of enhancing photosynthetic efficiency in C₃ plants by introducing CCM components from *Chlamydomonas*, we chose ten components that are considered essential or potentially important for CCM functionality (Table 1). These included intracellular components that mediate the transport and conversion of Ci from the external environment to the active sites of RuBisCO. LCI1 and HLA3 are putative Ci transporters reportedly in the plasma membrane; LCIA, CCP1 and CCP2 are putative chloroplast envelope Ci transporters, although only the former has been localized *in vivo*. CAH1, CAH3 and CAH6 are carbonic

anhydrases, thought to be located in the periplasmic space (CAH1), the thylakoid lumen (CAH3) and the chloroplast stroma (CAH6). LCIB and LCIC are proteins of unclear function that are reported to surround the pyrenoid when CO₂ becomes limiting.

We first expressed fluorescently tagged versions of all these proteins in *Chlamydomonas* cells to obtain definitive information about their locations and then expressed them transiently in tobacco (*Nicotiana benthamiana* L.) leaves. With one exception, proteins localized to identical compartments in the two organisms. Subsequent analyses focussed on the putative Ci transporters LCIA and HLA3, which have been shown to cooperatively drive bicarbonate uptake from the extracellular environment to the chloroplast stroma (Yamano *et al.*, 2015). We showed that both function as Ci transporters when expressed in the outer membrane of *Xenopus* oocytes. We also expressed these proteins stably in transgenic *Arabidopsis* plants, which grew as well as wild-type plants. Our results show that CCM components from *Chlamydomonas* can be expressed in appropriate locations in higher plant cells without compromising growth, although – consistent with modelling predictions – additional elements of the algal CCM will need to be co-expressed to achieve enhanced productivity.

Results

Subcellular localization of native CCM components in *Chlamydomonas*

The locations of CCM components in *Chlamydomonas* were investigated by transforming *Chlamydomonas* cells with constructs encoding these proteins fused to a fluorescent tag (Venus) at the C-terminus (Figures 1a and S1a). Full-length open reading frames were cloned from genomic DNA and constitutively expressed from the *PsaD* promoter. Most of the proteins had the subcellular locations expected from previous studies (Table 1). LCIA: Venus was confined to the chloroplast envelope; CAH3:




Figure 1 Expression of fluorescent-tagged CCM components in *Chlamydomonas* and tobacco. Expression of Venus-fused CCM components in *Chlamydomonas reinhardtii* (a). Expression in tobacco of GFP-fused CCM components from *Chlamydomonas* (b). Green and purple signals are Venus or GFP fluorescence and chlorophyll autofluorescence, respectively. Overlaid images of these signals are shown: overlaps are white. Scale bar = 5 μ m (all 5 μ m for *Chlamydomonas* images). For images of separate signals see Figure S1.

Venus, LCIB: Venus and LCIC: Venus were in the chloroplast, with LCIC: Venus and LCIB: Venus producing the distinctive circular pattern around the pyrenoid in CO_2 -starved cells. LCI1: Venus and HLA3: Venus were in the plasma membrane. In the absence of an available *Chlamydomonas* strain expressing tagged CAH1, we examined the location of the structurally related isozyme CAH2 (contiguous to CAH1 on chromosome 4, probably resulted from a gene duplication event, 91.8% identical amino acid sequences) (Fujiwara *et al.*, 1990). CAH2: Venus was at the cell periphery, consistent with the expected periplasmic locations of CAH1 and CAH2 (Ynalvez *et al.*, 2008). Unexpectedly, signals for CCP1: Venus and CCP2: Venus showed punctate subcellular localization, consistent with location in mitochondria. Furthermore, the location of fluorescence for CCP1: Venus and CCP2: Venus overlapped with that of a mitochondrial marker dye (Mitotracker Red CMXRos), indicating that both of these putative Ci transporters were located in mitochondria (Figure S2). There were no obvious signals for CAH6: Venus inside the *Chlamydomonas* cell (not shown).

Chlamydomonas CCM proteins can be expressed in tobacco leaves

The CCM components localized in *Chlamydomonas* cells were selected for expression in tobacco leaves. Binary expression vectors carrying each CCM gene individually were generated by PCR amplification of cDNA and subsequent Gateway cloning (Karimi *et al.*, 2002). Gene expression in tobacco was under the control of the constitutive 35S promoter and nopaline synthase (nos) terminator. Stop codons were removed to allow in-frame C-terminal fusion to a sequence encoding GFP. For CAH1, the N-terminal sequence (17 aa) was replaced with a leader sequence (22 aa) from tobacco, as described by Roberts and Spalding (1995), to facilitate processing and secretion to the apoplast.

The GFP-fused CCM components were expressed transiently in tobacco leaves by agro-infiltration. The locations of eight of the components were consistent with the demonstrated location in *Chlamydomonas* (Figures 1b and S1b). Fluorescent signals for LCIA: GFP were in the chloroplast envelope, and LCIB: GFP and




Figure 2 Co-expression of GFP-fused CCM components with a mCherry-fused plasma membrane transporter NPSN12 or a known mitochondrial marker (the targeting sequence of yeast cytochrome oxidase IV [COX4] fused to mCherry) in tobacco. Purple, green and cyan signals are chlorophyll autofluorescence, GFP and mCherry fluorescence, respectively. Overlaid images of these signals are shown: overlaps of GFP and mCherry are pale green. PM, plasma membrane; MT, mitochondria. Scale bar = 10 μm.




Figure 3 Expression of GFP-fused CCM components carrying native Arabidopsis chloroplast transit peptides in tobacco. Green and purple signals are GFP fluorescence and chlorophyll autofluorescence, respectively. Overlaid images of these signals are shown: overlaps are white. 1A-TP, RuBisCO small subunit RBCS1A (AT1G67090) transit peptide; ABC-TP, ABC transporter ABC113 (AT1G65410) transit peptide; mCAH6, mature CAH6; mLCIA, mature LCIA. Main image scale bar = 10 μ m, inset image scale bar = 3 μ m. For images of separate signals see Figure S3.

LCIC: GFP signals were stromal. LC11: GFP, HLA3: GFP and CAH1: GFP were at the cell periphery. The location of fluorescence for LC11: GFP and HLA3: GFP overlapped with that of an integral plasma membrane transporter protein (NPSN12, AT1G48240) fused to mCherry (Geldner *et al.*, 2009), indicating that both of these putative Ci transporters were associated with the plasma membrane (Figure 2). Although co-expression of CAH1: GFP with NPSN12: mCherry indicated that CAH1 was located at the cell periphery, we could not resolve whether the enzyme, which is periplasmic in *Chlamydomonas*, was expressed discretely in the intercellular space. The fluorescence signals for CCP1: GFP and CCP2: GFP appeared predominantly in numerous discrete structures much smaller than chloroplasts. This distribution is consistent with the locations of these proteins in mitochondria, as was the case in *Chlamydomonas* (Figure 1a). Furthermore, fluorescence signals for CCP1: GFP and CCP2: GFP overlapped with that of a mitochondrial marker (the targeting sequence of yeast cytochrome oxidase IV [COX4] fused to mCherry [Nelson *et al.*, 2007]), indicating that both of these putative Ci transporters were associated with the mitochondria (Figure 2).

The fluorescent signal for CAH3: GFP was present in both cytosol and chloroplasts when this fusion protein was expressed in tobacco. In contrast, CAH6: GFP appeared to be located exclusively in the cytosol (Figure 1b). Our own and previous data suggest that CAH3 is chloroplastic in *Chlamydomonas* and probably contained within the thylakoid lumen where it is suggested to play a pivotal role in the supply of CO₂ to RuBisCO (Figure 1) (Blanco-Rivero *et al.*, 2012; Duanmu *et al.*, 2009b; Karlsson *et al.*, 1998; Sinetova *et al.*, 2012). CAH6 has a predicted chloroplast transit peptide (TP), and there is some evidence for a chloroplast location in *Chlamydomonas* (Mitra *et al.*, 2004). However, our CAH6: Venus protein did not have a clear intracellular location. Thus, CAH3 appears to be mistargeted when expressed in tobacco, and there is insufficient information on CAH6 to determine whether this protein is correctly localized in tobacco.

Modification of target peptides for direction of CCM components to tobacco chloroplasts

We tested whether plasma membrane localized Ci transporters from *Chlamydomonas* could be retargeted to the chloroplast

envelope in tobacco leaves. Screening of the plant membrane protein database (Aramemnon, <http://aramemnon.botanik.uni-koeln.de/>) identified a chloroplast envelope transporter belonging to the ATP-binding cassette (ABC) superfamily: ABC113 of Arabidopsis (AT1G65410). The predicted N-terminal TP of ABC113 (ABC-TP, 60 aa) was attached to the N-termini of HLA3 and LC11. As controls, we used constructs encoding a *Chlamydomonas* protein: GFP fusion already shown to localize to the tobacco chloroplast envelope (LCIA, see above) that were modified either to remove the TP or to replace it with the ABC-TP.

When expressed transiently in tobacco, LCIA: GFP lacking the predicted native LCIA-TP (73 aa) (Miura *et al.*, 2004) localized to the cytosol instead of the chloroplast (Figures 3 and S3). Chloroplast envelope targeting was recovered with a N-terminal ABC-TP, similarly to full-length LCIA: GFP. This result showed that the ABC-TP could target some transporter proteins to the chloroplast envelope. However, the addition of the ABC-TP to LC11: GFP or to HLA3: GFP was not sufficient to retarget these two proteins to the chloroplast envelope. The fluorescence signal for ABC-TP: LC11: GFP indicated that the protein was in the chloroplast stroma, whereas the signal for ABC-TP: HLA3: GFP was in the cytosol.

Having found that CAH6: GFP did not localize to the chloroplast stroma, we replaced the native TP with the TP of the Arabidopsis RuBisCO small subunit, RBCS1A (AT1G67090). By placing the first 80 aa of RBCS1A (1A-TP) upstream of the mature CAH6 (mCAH6) to generate 1A-TP: mCAH6: GFP, the CAH6: GFP protein was retargeted to the chloroplast stroma (Figure 3). Thus, 1A-TP is a suitable sequence to direct the localization of soluble *Chlamydomonas* proteins to the stroma.

LCIA and HLA3 are located in the plasma membrane in *Xenopus* oocytes and increase Ci uptake rates

To investigate the putative function of LCIA and HLA3 as transmembrane Ci transporters, we expressed these proteins in *Xenopus* oocytes, with or without N-terminal GFP fusions. Oocytes injected with mRNA of either mature LCIA: GFP (mLCIA; lacking the N-terminal TP) or HLA3: GFP displayed a fluorescent signal on the cell surface after 3 d, indicating protein expression and incorporation into the plasma membrane (Figure 4a). mRNA-injected cells were assayed for Ci uptake using H¹⁴CO₃⁻ as a




Figure 4 Chlamydomonas CCM components LCIA and HLA3 facilitate increased accumulation of inorganic carbon in *Xenopus* oocytes. Confocal images of oocytes expressing GFP fused to mature LCIA (LCIA lacking a chloroplast transit peptide, mLCIA) or HLA3 3 d after injection (a). ^{14}C accumulation in oocytes expressing mLCIA or HLA3 either untagged or fused to GFP following 10-min incubation in MBS containing 0.12 mM $\text{NaH}^{14}\text{CO}_3$ (b). Values are means of measurements on 20 oocytes; bars are means \pm standard error (SE). Letters above the bars indicate a difference or between values; where a, b and c indicate significant difference ($P < 0.05$) as determined by analysis of variance (ANOVA) followed by Tukey's honestly significant difference (HSD) tests.

tracer. Oocytes transformed with mLCIA: GFP or HLA3: GFP accumulated 2.0- and 2.7-fold more ^{14}C than water-injected controls, respectively (Figure 4b). The presence of a GFP-tag had no adverse effect on $\text{H}^{14}\text{CO}_3^-$ uptake.

LCIA and HLA3 express in appropriate locations in *Arabidopsis* leaf cells following stable transformation

Following transformation by floral dip, three *Arabidopsis thaliana* (ecotype Columbia; Col-0) homozygous T3 lines with stable expression of either LCIA: GFP or HLA3: GFP were selected for further study. Both fusion proteins resulted in fluorescent signals in the same subcellular locations as in tobacco leaves (Figures 5a and S4). Leaf proteins were separated on SDS-PAGE and probed after blotting with a commercial antibody raised against GFP. Polypeptides corresponding to 54 kDa for LCIA: GFP and 170 kDa for HLA3: GFP were resolved (Figure 5b). These masses are consistent with those expected for GFP (27 kDa) fusions of LCIA after TP cleavage (27.5 kDa) and HLA3 (147 kDa) (Yamano *et al.*, 2015).

Transgenic *Arabidopsis* have normal growth and photosynthetic characteristics

Wild-type plants and LCIA: GFP- or HLA3: GFP-expressing lines were grown together under ambient CO_2 (ca. 400 $\mu\text{mol/mol}$) and

a light intensity of 100 $\mu\text{mol photons/m}^2/\text{s}$ (Figure 6a). Growth rate was compared by measuring rosette expansion, fresh weight and dry weight. No differences were observed between wild-type and transgenic lines. Furthermore, there was no difference in leaf chlorophyll content (Figure S5). To investigate whether the presence of algal C_i transporters affected growth rate under conditions in which chloroplastic CO_2 concentration is expected to be a major limitation on photosynthesis, plants were grown under low CO_2 (250 $\mu\text{mol/mol}$) and high light (350 $\mu\text{mol photons/m}^2/\text{s}$) (Figure 6b). All genotypes had higher growth rates and lower specific leaf areas under these conditions (Figure S6), but again, there were no significant differences in growth between wild-type and transgenic lines.

We checked whether expression of LCIA: GFP or HLA3: GFP affected the steady-state rate of photosynthetic CO_2 assimilation at ambient CO_2 (A_n) (Table 2). We also measured the relationship between the rate of photosynthesis and the CO_2 concentration in the leaf substomatal cavity (A/C_i , Figure 7) and used this to infer other photosynthetic parameters including the diffusion of CO_2 from substomatal cavity to chloroplast (mesophyll conductance, g_m) (Griffiths and Helliker, 2013) and the maximum RuBisCO carboxylation rate ($V_{c,\text{max}}$). There were no significant differences in any of the parameters between transgenic lines expressing LCIA: GFP or HLA3: GFP and wild-type plants.

Discussion

The introduction of a microbial eukaryotic CCM into crop plants will require robust methods to ensure that the proteins of interest perform their intended function in the foreign host system. To that effect, we developed a pipeline to (i) determine the native subcellular location in *Chlamydomonas*, (ii) test the targeting efficiency of constructs by transient expression in tobacco and (iii) provide a platform for physiological characterization of transgenic lines by stably transforming *Arabidopsis*. Through this approach, we showed that eight CCM components from *Chlamydomonas* can be successfully transferred into appropriate locations in leaves of higher plants. Confirmation of subcellular localization of CCM components is an important step in systematic introduction of a fully operational CCM into higher plants (Pengelly *et al.*, 2014). For example, successful expression and assembly of components of the cyanobacterial carboxysome microcompartment in tobacco chloroplasts will help to guide future attempts to assemble the cyanobacterial CCM into higher plants (Lin *et al.*, 2014a,b). In contrast, transfer of *ictB*, a putative cyanobacterial CCM component, into *Arabidopsis* and tobacco plants has resulted in enhanced biomass accumulation (Liemann-Hurwitz *et al.*, 2003; Simkin *et al.*, 2015), but lack of information on *ictB* protein structure, function and cellular localization in transgenic plants limits our understanding of this growth response (Simkin *et al.*, 2015).

The need to establish robustly the native location of CCM components, prior to incorporation into higher plants, is further supported by our work on two putative chloroplast membrane transporters. The suggestion that CCP1 and CCP2 are in the chloroplast envelope was based on their enrichment in chloroplast preparations and in preparations of envelope membranes (Mason *et al.*, 1990; Ramazanov *et al.*, 1993). The possibility of a mitochondrial location was not explicitly examined in these studies. Here, our robust co-expression approach revealed that tagged CCP1/2 proteins were associated with mitochondria rather than the chloroplast envelope in both *Chlamydomonas*




Figure 5 Stable expression of LCIA:GFP and HLA3:GFP in Arabidopsis. Representative confocal images of LCIA and HLA3 fused to GFP (a). Green and purple signals are GFP fluorescence and chlorophyll autofluorescence, respectively. Overlaid images of these signals are shown: overlaps are white. Scale bar = 10 µm. For images of separate signals see Figure S4. Immunoblots of rosette extracts (10 µg protein) from LCIA:GFP- and HLA3:GFP-expressing lines probed with an antibody against GFP (b). LCIA:GFP is present in three separate homozygous T3 insertion lines (LCIA:GFP₁₋₃), but not in segregating wild-type lines. HLA3:GFP is visible in HLA3:GFP₁₋₃ but not in the segregating wild-type for HLA3:GFP₁ or a wild-type equivalent for HLA3:GFP₂ and HLA3:GFP₃. LCIA:GFP and HLA3:GFP have approximate masses of 54 and 170 kDa, respectively (arrow). Ponceau stains of each blot (right) show the band attributable to the RuBisCO large subunit (RbCL, 55 kDa) as a loading control.

and tobacco (Figures 1 and 2; Figures S1 and S2). This location is also consistent with the high degree of similarity between CCP1/2 and the mitochondrial carrier protein superfamily (Pollock *et al.*, 2004; Pfam 00153; KOG0758). The expression of CCP1 and CCP2 is strongly induced by low CO₂, both genes are under the control of the CCM 'master switch' *Cia5/Ccm1*, and reducing CCP1/2 expression (through an RNAi approach) resulted in slower growth rates at low CO₂. However, cells with reduced CCP1/2 proteins did not lose CCM or photosynthetic capacity at low CO₂ (Pollock *et al.*, 2004). Taken together, these results suggest that transport across mitochondrial membranes plays an important role in coordinating the CCM with growth-related metabolism. In *Chlamydomonas*, mitochondria are known to relocate from a central position within the cup of the chloroplast to a peripheral position close to the plasma membrane when cells grown in high CO₂ are exposed to limiting CO₂ (Geraghty and Spalding, 1996). It has been suggested that mitochondrial relocation may be important either for energization of plasma membrane Ci transporters (to date, plasma membrane HLA3 is the only ATP-dependent Ci transporter candidate) or for scavenging glycolate produced in photorespiration during acclimation to limiting CO₂ (Spalding, 2009). A better understanding of the role of CCP1/2 in *Chlamydomonas* will inform future engineering strategies for higher plants.

The location of the carbonic anhydrases CAH3 and CAH6 in tobacco also differed from predicted results (Figure 2). The partial cytosolic localization of CAH3 is likely an effect of the failure of the higher plant import machinery to recognize the dual signalling peptide of this protein. In *Chlamydomonas* (Figure 1), results were consistent with location of the protein in the thylakoid lumen (Duanmu *et al.*, 2009b; Sinetova *et al.*, 2012). In particular, the presence of fluorescence in the centre of the

pyrenoid is consistent with the suggestion that CAH3 is concentrated in the transpyrenoidal thylakoid tubules following low CO₂-induced phosphorylation (Figure 1) (Blanco-Rivero *et al.*, 2012). In contrast, there is no clear evidence of a role for CAH6 in the *Chlamydomonas* CCM, and very little is known about its function. CAH6 represents the only apparent stromal carbonic anhydrase activity and has been adopted into the current CCM model as a speculative mechanism for Ci uptake and/or recovery of CO₂ escaping from the pyrenoid (Wang *et al.*, 2015; Yamano *et al.*, 2010). Evidence for chloroplast location in *Chlamydomonas* is based on a predicted stroma-signalling peptide and on immunogold labelling (Mitra *et al.*, 2004), but we did not observe an intracellular location for a CAH6: Venus protein.

Our attempts to relocate CCM components in tobacco cells by modifying targeting sequences were successful for some components. CAH6 could be redirected from the cytosol to the chloroplast stroma in tobacco by removal of the predicted TP and fusion to the TP of the Arabidopsis RuBisCO small subunit 1A, 1A-TP (Figure 3). The plasma membrane protein LC11 could be relocated to the chloroplast by fusion with the TP of an Arabidopsis chloroplast envelope transporter, and the same TP could substitute for the native TP of the *Chlamydomonas* chloroplast envelope protein LCIA. However, this TP did not redirect the plasma membrane protein HLA3 to the chloroplast envelope. Moving larger transmembrane proteins like HLA3 to the chloroplast may require further modifications to remove potential competing signal motifs and to ensure correct orientation in the case of directional channels.

The putative Ci transporters HLA3 and LCIA localized to the plasma membrane and chloroplast envelope, respectively, in *Chlamydomonas*, tobacco and Arabidopsis (Figures 1 and 5). Recent work has confirmed these locations in *Chlamydomonas*





Figure 6 Growth of phenotypes in different environmental conditions of transgenic Arabidopsis plants expressing LCIA or HLA3. Plants were grown under ambient CO₂ (ca. 400 μmol/mol) and 100 μmol photons/m²/s (a) or low CO₂ (250 μmol/mol) and 350 μmol photons/m²/s (b). Growth rates (1st and 3rd row) and fresh weight (FW) and dry weight (DW) (2nd and 4th row) are shown for LCIA and HLA3, respectively. HLA3 transgenic lines had a lower FW and DW compared to LCIA when grown under ambient CO₂, as plants were harvested slightly earlier (at 29 days vs 31 days). All plants grown under low CO₂ were harvested at 30 days. Values are the means ± SE of measurements made on 24 rosettes.

Table 2 Photosynthetic parameters determined from gas exchange analysis of LCIA or HLA3 transgenic plants. Values are the mean \pm SE of measurements made on four leaves, each from a different plant (as shown in Figure 7)

	Wild type	HLA3: GFP ₁	HLA3: GFP ₂	HLA3: GFP ₃	wild type	LCIA: GFP ₁	LCIA: GFP ₂	LCIA: GFP ₃
A_n ($\mu\text{mol CO}_2/\text{m}^2/\text{s}$)	9.4 \pm 0.6	11 \pm 0.9	10 \pm 0.8	9.9 \pm 0.8	9.7 \pm 0.2	9.4 \pm 0.5	9.7 \pm 1.2	8.8 \pm 0.2
g_s (mol CO ₂ /m ² /s)	0.21 \pm 0.04	0.29 \pm 0.03	0.21 \pm 0.03	0.25 \pm 0.04	0.26 \pm 0.03	0.24 \pm 0.04	0.25 \pm 0.05	0.24 \pm 0.01
g_m (mol CO ₂ /m ² /s)	0.047 \pm 0.003	0.048 \pm 0.005	0.047 \pm 0.003	0.044 \pm 0.005	0.04 \pm 0.001	0.038 \pm 0.002	0.04 \pm 0.004	0.034 \pm 0.001
$V_{c,\text{max}}$ ($\mu\text{mol CO}_2/\text{m}^2/\text{s}$)	30 \pm 3.2	30.4 \pm 2.4	29.3 \pm 1.8	27.3 \pm 2.2	28.7 \pm 1.1	29.2 \pm 2.8	31.4 \pm 2.3	27 \pm 0.7
J_{max} ($\mu\text{mol e}^-/\text{m}^2/\text{s}$)	64.5 \pm 4.8	71.6 \pm 5.6	67 \pm 4.7	64.3 \pm 5.2	64.5 \pm 1.9	65.1 \pm 4.6	69.5 \pm 6.6	61.5 \pm 1.1
Γ ($\mu\text{mol CO}_2/\text{mol}$)	39.5 \pm 2.5	44 \pm 2.6	37.9 \pm 1.4	36.1 \pm 2.6	33.2 \pm 0.6	31.5 \pm 2.4	32.9 \pm 0.9	29.1 \pm 1
Initial slope (A_n/C_i)	0.044 \pm 0.002	0.046 \pm 0.004	0.044 \pm 0.003	0.043 \pm 0.004	0.038 \pm 0.001	0.035 \pm 0.002	0.038 \pm 0.005	0.036 \pm 0.001

A_n , net photosynthesis at ambient CO₂; g_s , stomatal conductance to CO₂; g_m , mesophyll conductance to CO₂; $V_{c,\text{max}}$, maximum velocity of RuBisCO carboxylation; J_{max} , maximum capacity of electron transport; Γ , CO₂ compensation point.

ANOVA revealed that there were no statistically significant differences between samples ($P < 0.05$).

**Figure 7** Photosynthetic responses of transgenic plants. Photosynthetic rates were determined as a function of increasing substomatal CO₂ concentrations (A/C_i) at saturating light levels (1500 $\mu\text{mol photons}/\text{m}^2/\text{s}$). Each curve represents the means \pm SE of values from four leaves, each on a different plant.

(Yamano *et al.*, 2015). We showed that both proteins can facilitate uptake of C_i into *Xenopus* oocytes, consistent with previous demonstrations that LCIA can facilitate C_i movement across membranes (Mariscal *et al.*, 2006) and that HLA3 overexpression contributes to increased C_i uptake in *Chlamydomonas* (Gao *et al.*, 2015; Yamano *et al.*, 2015). Further kinetic analyses will be required to determine the mode of action of these proteins (i.e. active vs. passive transport). This information will be crucial for modelling approaches and rational engineering strategies. For example, rational approaches for engineering the cyanobacterial CCM into higher plants are based on a good understanding of the catalytic characteristics of cyanobacterial C_i transporters BicA and SbtA (Du *et al.*, 2014; McGrath and Long, 2014; Price *et al.*, 2011, 2013).

The addition of LCIA or HLA3 did not confer a growth advantage to *Arabidopsis* plants (Figures 6 and 7, Table 2). This is not entirely unexpected, as biophysical CCMs, in both algae and cyanobacteria, require additional features to function, including a

microcompartment containing RuBisCO and additional components to reduce C_i leakage (Badger *et al.*, 1998; Price *et al.*, 2013; Yamano *et al.*, 2010). Furthermore, it is likely that native carbonic anhydrase activity, particularly in the stroma, would hinder C_i accumulation. Biophysical CCMs rely on the capacity to accumulate bicarbonate; thus, specific localization and control of carbonic anhydrase activity appear to play an important role in functionality. For example, ectopic expression of carbonic anhydrase within the cytoplasm of cyanobacterial cells leads to a debilitating leakage of C_i due to rapid equilibration between bicarbonate and CO₂ in the cytosol (Price and Badger, 1989). Predictive models indicate that the removal of native stromal carbonic anhydrases is a key target for introducing biophysical CCMs into higher plants (McGrath and Long, 2014; Price *et al.*, 2013). Our results do demonstrate that higher plant cells can express C_i transporters from *Chlamydomonas* in appropriate locations, without deleterious effect. Both LCIA and HLA3 showed appropriate transmembrane integration and increased

Ci uptake when expressed in *Xenopus* oocytes (Figures 1, 4 and 5). Furthermore, we were able to detect both proteins in transgenic *Arabidopsis* leaves (Figure 5). Together, these data suggest that LCIA and HLA3 were active, but that activity was not sufficient to drive up CO₂ levels around RuBisCO and hence improve the rate of CO₂ assimilation.

Recent work has shown that expression and function of HLA3 and LCIA may be closely coordinated in *Chlamydomonas* (Yamano *et al.*, 2015). Enhanced Ci uptake was observed when both proteins were overexpressed in wild-type *Chlamydomonas* at high CO₂. However, negligible changes in photosynthesis and Ci uptake were observed when either HLA3 or LCIA was overexpressed, indicating that, in the absence of other CCM components or a microcompartment such as a pyrenoid, these proteins are not able to enhance chloroplastic CO₂ concentrations (Gao *et al.*, 2015; Yamano *et al.*, 2015). These data suggest that subsequent engineering strategies in higher plants should focus on co-expression of HLA3 and LCIA. Additional modifications will probably be required to establish a functional biophysical CCM in higher plants, including the removal of native carbonic anhydrases, which would otherwise short circuit active uptake mechanisms and eliminate any stromal pool of bicarbonate (Price *et al.*, 2013). The next challenge will be to stack key CCM components, and develop a strategy to retarget stromal RuBisCO to a chloroplast microcompartment, a predicted requirement for an algal-type CCM (Badger *et al.*, 1998).

Experimental procedures

Plant material and growth conditions

Mutant and transgenic plants of *Arabidopsis* (*Arabidopsis thaliana*) were in the Col-0 wild-type background. *Arabidopsis* seeds were sown on compost, stratified for 3 days at 4 °C and grown at 20 °C, under ambient CO₂ (ca. 400 µmol/mol), at 70% relative humidity and under 100 µmol photons/m²/s in 12-h light/12-h dark cycles, unless otherwise stated.

For analyses of transgenic lines, homozygous insertion or wild-type out-segregant lines (T3) were compared. Where wild-type out-segregants were not available, homozygous insertion lines were compared with Col-0 plants from seed stocks of the same age generated under similar conditions. Tobacco plants (*Nicotiana benthamiana* L.) were cultivated under glass house conditions (minimum 20 °C, natural light supplemented to give at least 12-h light). Venus-tagged proteins were expressed in wild-type *Chlamydomonas* strain cMJ030 (CC-4533) (Zhang *et al.*, 2014). Cells were maintained in constant low light (~10 µmol photons/m²/s) at RT on 1.5% (w/v) agar plates containing Tris–acetate–phosphate (TAP) (Kropat *et al.*, 2011). For imaging, cells were grown in liquid TAP media to a concentration of 10⁶ cells/mL, pelleted by centrifugation (1000 g, 4 min), resuspended in Tris–phosphate (T-P) minimal media (Kropat *et al.*, 2011) and grown for 24 h in ambient CO₂ before imaging.

Cloning and expression of CCM components in *Chlamydomonas*

The open reading frames (ORFs) of *Chlamydomonas* genes were expressed in frame with Venus from the *PsaD* promoter using the pLM005 vector. ORFs were amplified from genomic DNA using Phusion Hotstart II polymerase (Thermo Fisher Scientific, www.thermofisher.com) with the respective oligos in Table S1. HpaI-cut pLM005 vector and PCR products were gel purified and assembled by Gibson assembly (Gibson *et al.*, 2009). Due to the

large gene length of *HLA3*, it was cloned in two fragments then assembled in the pLM005 vector by Gibson assembly. The pLM005 vector contains the AphVIII gene for paromomycin resistance in *Chlamydomonas* and ampicillin resistance for bacterial selection. All construct junctions were verified by Sanger sequencing. Constructs were transformed into *Chlamydomonas* by electroporation as in Zhang *et al.* (2014). Briefly, 250 µL of 2 × 10⁸ cells/mL was transformed with 14.5 ng/kbp of EcoRV-cut plasmid at 16 °C. Cells were spread on 86 mL TAP agar plates containing paromomycin (20 µg/mL) and kept in low light (~10 µmol photons/m²/s) until colonies were ~2–3 mm in diameter. Plates were screened for fluorescent colonies using a Typhoon TRIO fluorescence scanner (GE Healthcare, www.gelifesciences.com) with excitation/emission wavelengths 532 nm/520–555 nm for Venus and 633 nm/630–670 nm for chlorophyll autofluorescence.

Cloning and expression of CCM components in tobacco and *Arabidopsis*

Genes were cloned from cDNA derived from *Chlamydomonas* (strain CC-4886, *Chlamydomonas* Resource Center). Primers were designed from sequences available on Phytozome v10.2 (*Chlamydomonas reinhardtii* v5.5 [Augustus u11.6], http://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org_Creinhardtii) (see Table S1 for oligo details). Gene sequences for *LCIA* and *HLA3* were codon-optimized for expression in higher plants and synthesized *de novo* (DNA2.0, CA, USA) (Figure S7), then cloned into Gateway entry vectors (pCR[®]8/GW/TOPO[®]TA Cloning[®] Kit) using Platinum[®] Taq DNA Polymerase High Fidelity according to the manufacturer's instructions (Invitrogen[™] Life Technologies, www.lifetechnologies.com) and subsequently cloned into the destination binary vectors pK7FWG2.0 (Karimi *et al.*, 2002) or pGWB5 (Nakagawa *et al.*, 2009). Gibson assembly was used to generate transit peptide gene fusions. Binary vectors were transformed into *Agrobacterium tumefaciens* (AGL1) for transient gene expression in tobacco leaves (Schöb *et al.*, 1997) or stable insertion in *Arabidopsis* plants by floral dipping (Clough and Bent, 1998). Co-expression studies were performed with the WAVE131 vector of the 'wave' marker set (Geldner *et al.*, 2009) and the mt-rb vector (Nelson *et al.*, 2007) for plasma membrane and mitochondria localization, respectively.

DNA extraction, PCR and protein analysis

For screening transgenic *Arabidopsis* lines, genomic DNA was extracted from mature, nonflowering rosettes of T1 plants as described in Li and Chory (Li and Chory, 1998). PCRs were performed as in McCormick and Kruger (2015). Where possible, the location of gene inserts was confirmed by TAIL PCR as described by Liu *et al.* (1995). Homozygous insertion lines were identified in the T2 generation either by PCR or by seedling segregation ratios on kanamycin-containing Murashige and Skoog (MS) medium (0.5x) plates.

Relative levels of LCIA: GFP and HLA3: GFP proteins in leaves were confirmed by immunoblot. Approximately 10 µg protein from whole 28-d-old rosettes (100 mg fresh weight) was fractionated by SDS-PAGE on a 10% (w/v) acrylamide: bisacrylamide (40:1) gel transferred to PVDF membrane, probed with mouse anti-GFP IgG_{2a} at 1:1,000 dilution (Santa Cruz, <http://www.scbt.com/>) and visualized using an HRP-conjugated goat anti-mouse IgG_{2a} at 1:50 000 dilution. HRP activity was detected using Supersignal Ultra (Pierce, www.piercenet.com) according to the manufacturer's instructions.

Oocyte expression and bicarbonate uptake assays

Synthesized gene sequences for mature *LCIA* (mLCIA) or *HLA3* lacking a stop codon were cloned into the expression vector Vivid Colors™ pcDNA™ 6.2/EmGFP (Invitrogen™ Life Technologies) to generate mLCIA- or HLA3-GFP fusions. For LCIA, the N-terminal transit sequence peptide (73 aa) was removed and replaced with the sequence 'GACATG' to add a Kozak sequence and new start codon. For HLA3, 'GAC' was added immediately upstream of the start codon. To generate equivalent vectors lacking a fluorescent tag, Gibson assembly was used to add a stop codon to mLCIA or HLA3 and remove the GFP sequence (720 bp). Plasmids were linearized by *AvrII* or *StuI* (Roche, www.roche.co.uk) and capped mRNA was synthesized using the mMESSAGE mMACHINE® T7 Transcription Kit (Ambion®; Life Technologies) according to the manufacturer's instructions. Mature LCIA or HLA3 mRNA was expressed in oocytes as described by Feng *et al.* (2013). *Xenopus* oocytes were injected with 50 nL of mRNA (1 µg/µL) or diethyl pyrocarbonate (DEPC)-treated water as a control. Bicarbonate uptake assays and confocal imaging were performed 3 d after injection in a protocol adapted from Mariscal *et al.* (2006). Oocytes were incubated in fresh MBS media (88 mM NaCl, 1 mM KCl, 2.4 mM NaHCO₃, 0.71 mM CaCl₂, 0.82 mM MgSO₄ and 15 mM HEPES, pH 7.4), containing 0.12 mM NaHCO₃ (1.85 GBq/mol NaH¹⁴CO₃). After 10 min, the oocytes were washed three times with ice-cold MBS and lysed in 200 µL SDS (10% [w/v]), and the radioactivity retained in individual oocytes was measured.

Chlorophyll quantification

Chlorophyll was extracted from powdered leaf discs in ice-cold 80% (v/v) acetone and 10 mM Tris-HCl, and concentration was measured according to Porra *et al.* (1989).

Measurement of photosynthetic parameters

Gas exchange rates were determined using a LI-6400 portable infrared gas analyser (LI-COR Biosciences, <http://www.licor.com/>) on either the sixth or seventh leaf of 35- to 45-d-old mature, nonflowering rosettes grown in large pots to generate leaf area sufficient for gas exchange measurements (Flexas *et al.*, 2007). The response of net photosynthetic CO₂ assimilation (*A*) to substomatal CO₂ concentration (*C_i*) was measured by varying the external CO₂ concentration from 0 to 1000 µmol/mol under a constant photosynthetic active radiation of 1500 µmol photons/m²/s (provided by a red-blue light source attached to the leaf chamber). Gas exchange data were corrected for CO₂ diffusion as in Bellasio *et al.* (2015). Leaf temperature and chamber relative humidity were maintained at 21 °C and 70%, respectively. To calculate maximum carboxylation rate (*V_{c,max}*), maximum electron transport flow (*J_{max}*) and mesophyll conductance (*g_m*), the *A/C_i* data were fitted to the *C₃* photosynthesis model (Farquhar *et al.*, 1980) with modifications to include estimations for *g_m* as described by Ethier and Livingston (2004).

Confocal laser scanning microscopy

A Leica TCS SP2 laser scanning confocal microscope (Leica Microsystems) with a water immersion objective lens (HCX IRAPO 25.0x0.95) was used for imaging leaves and oocytes. Excitation/emission wavelengths were 488 nm/500–530 nm for GFP, 543 nm/590–620 nm for mCherry and 488 nm/680–750 nm for chlorophyll autofluorescence. Images were acquired using Leica LAS AF software (<http://www.leica-microsystems.com/>). Prior to imaging Venus-tagged proteins in *Chlamydomonas*,

15 µL of cells were added to a well of a 96-well optical plate (Brooks Life Science Systems, <http://www.brooks.com>) and covered with 150 µL of 1.5% low melting point agarose containing T-P (~35 °C). For mitochondria staining, CCP1: Venus- and CCP2: Venus-expressing lines were grown in liquid T-P media with paromomycin (2 µg/mL) to a concentration of 2–4 × 10⁶ cells/mL. Cultures were incubated with MitoTracker Red CMXRos (Thermo Fisher Scientific) to a final concentration of 1 µM for 10 min, then spotted on a polylysine-coated slide for imaging. Cells were imaged using a custom adapted confocal microscope (Leica DMI6000) with settings at 514 nm/532–555 nm for Venus, 561 nm/573–637 nm for staining by Mitotracker Red CMXRos and 561 nm/665–705 nm for chlorophyll autofluorescence. Images were analysed using Fiji software (<http://fiji.sc/Fiji>).

Statistical analysis

Variations in response between genotypes were assessed by either analysis of variance (ANOVA) or Student's *t*-tests followed by Tukey's honest significant difference (HSD) post hoc test (SPSS Statistics 18, <http://www.ibm.com/>). Differences for which *P* < 0.05 are considered significant.

Acknowledgements

We thank Livia Scheunemann, Xiaorong Fan, Paloma Menguer and Anthony Miller (John Innes Centre) for advice and help with oocyte experiments, Mark Fricker (University of Oxford) for use of LeafLab software and Fani Ntana (University of Edinburgh) for technical contributions. AJM, DF and AMS were funded by grants from the Biotechnology and Biological Sciences Research Council (BBSRC), UK: BB/I024453/1 and Institute Strategic Programme Grant BB/J004561/1 to the John Innes Centre. NA was funded by BBSRC: BB/M006468/1, HG and MM were funded by BBSRC: BB/M007693/1. LM and MJ were funded by US National Science Foundation (NSF): 1359682.

Conflict of interest

The authors declare that they have no competing interests.

References

- Badger, M.R., Andrews, T.J., Whitney, S.M., Ludwig, M., Yellowlees, D.C., Leggat, W. and Price, G.D. (1998) The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae. *Can. J. Bot.* **76**, 1052–1071.
- Bellasio, C., Beerling, D.J. and Griffiths, H. (2015) An Excel tool for deriving key photosynthetic parameters from combined gas exchange and chlorophyll fluorescence: theory and practice. *Plant, Cell Environ.* doi: 10.1111/pce.12560.
- Blanco-Rivero, A., Shutova, T., Roman, M.J., Villarejo, A. and Martinez, F. (2012) Phosphorylation controls the localization and activation of the lumenal carbonic anhydrase in *Chlamydomonas reinhardtii*. *PLoS ONE*, **7**, e49063.
- Brueggeman, A.J., Gangadharaiah, D.S., Cserhati, M.F., Casero, D., Weeks, D.P. and Ladunga, I. (2012) Activation of the carbon concentrating mechanism by CO₂ deprivation coincides with massive transcriptional restructuring in *Chlamydomonas reinhardtii*. *Plant Cell*, **24**, 1860–1875.
- Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735–743.
- Du, J., Forster, B., Rourke, L., Howitt, S.M. and Price, G.D. (2014) Characterisation of cyanobacterial bicarbonate transporters in *E. coli* shows that SbtA homologs are functional in this heterologous expression system. *PLoS ONE*, **9**, e115905.

- Duanmu, D., Miller, A.R., Horken, K.M., Weeks, D.P. and Spalding, M.H. (2009a) Knockdown of limiting- CO_2 -induced gene HLA3 decreases HCO_3^- transport and photosynthetic C_i affinity in *Chlamydomonas reinhardtii*. *Proc. Natl Acad. Sci. USA*, **106**, 5990–5995.
- Duanmu, D., Wang, Y. and Spalding, M.H. (2009b) Thylakoid lumen carbonic anhydrase (CAH3) mutation suppresses air-Dier phenotype of LCIB mutant in *Chlamydomonas reinhardtii*. *Plant Physiol.* **149**, 929–937.
- Engel, B.D., Schaffer, M., Kuhn Cuellar, L., Villa, E., Plitzko, J.M. and Baumeister, W. (2015) Native architecture of the *Chlamydomonas* chloroplast revealed by in situ cryo-electron tomography. *eLife*, **4**, e04889.
- Ethier, G.J. and Livingston, N.J. (2004) On the need to incorporate sensitivity to CO_2 transfer conductance into the Farquhar-von Caemmerer-Berry leaf photosynthesis model. *Plant, Cell Environ.* **27**, 137–153.
- Fang, W., Si, Y., Douglass, S., Casero, D., Merchant, S.S., Pellegrini, M., Ladunga, I. et al. (2012) Transcriptome-wide changes in *Chlamydomonas reinhardtii* gene expression regulated by carbon dioxide and the CO_2 -concentrating mechanism regulator CIA5/CCM1. *Plant Cell*, **24**, 1876–1893.
- Farquhar, G.D., von Caemmerer, S. and Berry, J.A. (1980) A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species. *Planta*, **149**, 78–90.
- Feng, H., Xia, X., Fan, X., Xu, G. and Miller, A.J. (2013) Optimizing plant transporter expression in *Xenopus* oocytes. *Plant Methods*, **9**, 48.
- Flexas, J., Ortuno, M.F., Ribas-Carbo, M., Diaz-Espejo, A., Florez-Sarasa, I.D. and Medrano, H. (2007) Mesophyll conductance to CO_2 in *Arabidopsis thaliana*. *New Phytol.* **175**, 501–511.
- Fujiwara, S., Fukuzawa, H., Tachiki, A. and Miyachi, S. (1990) Structure and differential expression of two genes encoding carbonic anhydrase in *Chlamydomonas reinhardtii*. *Proc. Natl Acad. Sci. USA*, **87**, 9779–9783.
- Gao, H., Wang, Y., Fei, X., Wright, D.A. and Spalding, M.H. (2015) Expression activation and functional analysis of HLA3, a putative inorganic carbon transporter in *Chlamydomonas reinhardtii*. *Plant J.* **82**, 1–11.
- Geldner, N., Dénervaud-Tendon, V., Hyman, D.L., Mayer, U., Stierhof, Y.-D. and Chory, J. (2009) Rapid, combinatorial analysis of membrane compartments in intact plants with a multicolor marker set. *Plant J.* **59**, 169–178.
- Genkov, T., Meyer, M., Griffiths, H. and Spreitzer, R.J. (2010) Functional hybrid Rubisco enzymes with plant small subunits and algal large subunits: engineered rbcS cDNA for expression in *Chlamydomonas*. *J. Biol. Chem.* **285**, 19833–19841.
- Geraghty, A.M. and Spalding, M.H. (1996) Molecular and structural changes in *Chlamydomonas* under limiting CO_2 (a possible mitochondrial role in adaptation). *Plant Physiol.* **111**, 1339–1347.
- Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A. 3rd and Smith, H.O. (2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods*, **6**, 343–345.
- Griffiths, H. and Helliker, B.R. (2013) Mesophyll conductance: internal insights of leaf carbon exchange. *Plant, Cell Environ.* **36**, 733–735.
- Karimi, M., Inze, D. and Depicker, A. (2002) Gateway vectors for *Agrobacterium*-mediated plant transformation. *Trends Plant Sci.* **7**, 193–195.
- Karlsson, J., Clarke, A.K., Chen, Z.Y., Huggins, S.Y., Park, Y.I., Husic, H.D., Moroney, J.V. et al. (1998) A novel alpha-type carbonic anhydrase associated with the thylakoid membrane in *Chlamydomonas reinhardtii* is required for growth at ambient CO_2 . *EMBO J.* **17**, 1208–1216.
- Kimpel, D.L., Togasaki, R.K. and Miyachi, S. (1983) Carbonic anhydrase in *Chlamydomonas reinhardtii* I. Localization. *Plant Cell Physiol.* **24**, 255–259.
- Kropat, J., Hong-Hermesdorf, A., Casero, D., Ent, P., Castruita, M., Pellegrini, M., Merchant, S.S. et al. (2011) A revised mineral nutrient supplement increases biomass and growth rate in *Chlamydomonas reinhardtii*. *Plant J.* **66**, 770–780.
- Li, J. and Chory, J. (1998) Preparation of DNA from *Arabidopsis*. *Methods Mol. Biol.* **82**, 55–60.
- Lieman-Hurwitz, J., Rachmilevitch, S., Mittler, R., Marcus, Y. and Kaplan, A. (2003) Enhanced photosynthesis and growth of transgenic plants that express ictB, a gene involved in HCO_3^- accumulation in cyanobacteria. *Plant Biotechnol. J.* **1**, 43–50.
- Lin, M.T., Occhialini, A., Andralojc, P.J., Devonshire, J., Hines, K.M., Parry, M.A.J. and Hanson, M.R. (2014a) β -Carboxysomal proteins assemble into highly organized structures in *Nicotiana* chloroplasts. *Plant J.* **79**, 1–12.
- Lin, M.T., Occhialini, A., Andralojc, P.J., Parry, M.A.J. and Hanson, M.R. (2014b) A faster Rubisco with potential to increase photosynthesis in crops. *Nature*, **513**, 547–550.
- Liu, Y.-G., Mitsukawa, N., Oosumi, T. and Whittier, R.F. (1995) Efficient isolation and mapping of *Arabidopsis thaliana* T-DNA insert junctions by thermal asymmetric interlaced PCR. *Plant J.* **8**, 457–463.
- Long, S.P., Zhu, X.G., Naidu, S.L. and Ort, D.R. (2006) Can improvement in photosynthesis increase crop yields? *Plant, Cell Environ.* **29**, 315–330.
- Long, S.P., Marshall-Colon, A. and Zhu, X.G. (2015) Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell*, **161**, 56–66.
- Mariscal, V., Moulin, P., Orsel, M., Miller, A.J., Fernandez, E. and Galvan, A. (2006) Differential regulation of the *Chlamydomonas* Nar1 gene family by carbon and nitrogen. *Protist*, **157**, 421–433.
- Markelova, A.G., Sinetova, M.P., Kupriyanova, E.V. and Pronina, N.A. (2009) Distribution and functional role of carbonic anhydrase Cah3 associated with thylakoid membranes in the chloroplast and pyrenoid of *Chlamydomonas reinhardtii*. *Russ. J. Plant Physiol.* **56**, 761–768.
- Mason, C.B., Manuel, L.J. and Moroney, J.V. (1990) A new chloroplast protein is induced by growth on low CO_2 in *Chlamydomonas reinhardtii*. *Plant Physiol.* **93**, 833–836.
- McCormick, A.J. and Kruger, N.J. (2015) Lack of fructose 2,6-bisphosphate compromises photosynthesis and growth in *Arabidopsis* in fluctuating environments. *Plant J.* **81**, 670–683.
- McGrath, J.M. and Long, S.P. (2014) Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis. *Plant Physiol.* **164**, 2247–2261.
- Meyer, M. and Griffiths, H. (2013) Origins and diversity of eukaryotic CO_2 -concentrating mechanisms: lessons for the future. *J. Exp. Bot.* **64**, 769–786.
- Meyer, M.T., Genkov, T., Skepper, J.N., Jouhet, J., Mitchell, M.C., Spreitzer, R.J. and Griffiths, H. (2012) Rubisco small-subunit alpha-helices control pyrenoid formation in *Chlamydomonas*. *Proc. Natl Acad. Sci. USA*, **109**, 19474–19479.
- Mitra, M., Lato, S.M., Ynalvez, R.A., Xiao, Y. and Moroney, J.V. (2004) Identification of a new chloroplast carbonic anhydrase in *Chlamydomonas reinhardtii*. *Plant Physiol.* **135**, 173–182.
- Miura, K., Yamano, T., Yoshioka, S., Kohinata, T., Inoue, Y., Taniguchi, F., Asamizu, E. et al. (2004) Expression profiling-based identification of CO_2 -responsive genes regulated by CCM1 controlling a carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Physiol.* **135**, 1595–1607.
- Morita, E., Abe, T., Tsuzuki, M., Fujiwara, S., Sato, N., Hirata, A., Sonoike, K. et al. (1998) Presence of the CO_2 -concentrating mechanism in some species of the pyrenoid-less free-living algal genus *Chloromonas* (Volvocales, Chlorophyta). *Planta*, **204**, 269–276.
- Moroney, J.V. and Tolbert, N.E. (1985) Inorganic carbon uptake by *Chlamydomonas reinhardtii*. *Plant Physiol.* **77**, 253–258.
- Nakagawa, T., Ishiguro, S. and Kimura, T. (2009) Gateway vectors for plant transformation. *Plant Biotechnol.* **26**, 275–284.
- Nelson, B.K., Cai, X. and Nebenführ, A. (2007) A multicolored set of *in vivo* organelle markers for co-localization studies in *Arabidopsis* and other plants. *Plant J.* **51**, 1126–1136.
- Ohnishi, N., Mukherjee, B., Tsujikawa, T., Yanase, M., Nakano, H., Moroney, J.V. and Fukuzawa, H. (2010) Expression of a low CO_2 -inducible protein, LC11, increases inorganic carbon uptake in the green alga *Chlamydomonas reinhardtii*. *Plant Cell*, **22**, 3105–3117.
- Parry, M.A., Andralojc, P.J., Scales, J.C., Salvucci, M.E., Carmo-Silva, A.E., Alonso, H. and Whitney, S.M. (2013) Rubisco activity and regulation as targets for crop improvement. *J. Exp. Bot.* **64**, 717–730.
- Pengelly, J.J., Forster, B., von Caemmerer, S., Badger, M.R., Price, G.D. and Whitney, S.M. (2014) Transplastomic integration of a cyanobacterial bicarbonate transporter into tobacco chloroplasts. *J. Exp. Bot.* **65**, 3071–3080.
- Pollock, S.V., Prout, D.L., Godfrey, A.C., Lemaire, S.D. and Moroney, J.V. (2004) The *Chlamydomonas reinhardtii* proteins Ccp1 and Ccp2 are required for long-term growth, but are not necessary for efficient photosynthesis, in a low- CO_2 environment. *Plant Mol. Biol.* **56**, 125–132.
- Porra, R.J., Thompson, W.A. and Kriedemann, P.E. (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the

- concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochim. Biophys. Acta*, **975**, 384–394.
- Price, G.D. and Badger, M.R. (1989) Expression of human carbonic anhydrase in the cyanobacterium *Synechococcus* PCC7942 creates a high CO₂-requiring phenotype - evidence for a central role for carboxysomes in the CO₂ concentrating mechanism. *Plant Physiol.* **91**, 505–513.
- Price, G.D., Badger, M.R. and von Caemmerer, S. (2011) The prospect of using cyanobacterial bicarbonate transporters to improve leaf photosynthesis in C3 crop plants. *Plant Physiol.* **155**, 20–26.
- Price, G.D., Pengelly, J.J., Forster, B., Du, J., Whitney, S.M., von Caemmerer, S., Badger, M.R. et al. (2013) The cyanobacterial CCM as a source of genes for improving photosynthetic CO₂ fixation in crop species. *J. Exp. Bot.* **64**, 753–768.
- Ramazanov, Z., Mason, C.B., Geraghty, A.M., Spalding, M.H. and Moroney, J.V. (1993) The Low CO₂-Inducible 36-kilodalton protein is localized to the chloroplast envelope of *Chlamydomonas reinhardtii*. *Plant Physiol.* **101**, 1195–1199.
- Roberts, C.S. and Spalding, M.H. (1995) Post-translational processing of the highly processed, secreted periplasmic carbonic anhydrase of *Chlamydomonas* is largely conserved in transgenic tobacco. *Plant Mol. Biol.* **29**, 303–315.
- Schöb, H., Kunz, C. and Meins, F. Jr. (1997) Silencing of transgenes introduced into leaves by agroinfiltration: a simple, rapid method for investigating sequence requirements for gene silencing. *Mol. Gen. Genet.* **256**, 581–585.
- Sharkey, T.D. (1988) Estimating the rate of photorespiration in leaves. *Physiol. Plant.* **73**, 147–152.
- Simkin, A.J., McAusland, L., Headland, L.R., Lawson, T. and Raines, C.A. (2015) Multigene manipulation of photosynthetic carbon assimilation increases CO₂ fixation and biomass yield in tobacco. *J. Exp. Bot.* **66**, 4075–4090.
- Sinetova, M.A., Kupriyanova, E.V., Markelova, A.G., Allakhverdiev, S.I. and Pronina, N.A. (2012) Identification and functional role of the carbonic anhydrase Cah3 in thylakoid membranes of pyrenoid of *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, **1817**, 1248–1255.
- Spalding, M.H. (2009) The CO₂-concentrating mechanism and carbon assimilation. In *The Chlamydomonas Sourcebook: Organellar and Metabolic Processes* (Harris, E.H. and Stern, D.B., eds), pp. 257–301. Oxford: Academic.
- Villarejo, A., Shutova, T., Moskvina, O., Forssen, M., Klimov, V.V. and Samuelsson, G. (2002) A photosystem II-associated carbonic anhydrase regulates the efficiency of photosynthetic oxygen evolution. *EMBO J.* **21**, 1930–1938.
- Wang, Y. and Spalding, M.H. (2014a) Acclimation to very low CO₂: contribution of limiting CO₂ inducible proteins, LCIB and LCIA, to inorganic carbon uptake in *Chlamydomonas reinhardtii*. *Plant Physiol.* **166**, 2040–2050.
- Wang, Y. and Spalding, M.H. (2014b) LCIB in the *Chlamydomonas* CO₂-concentrating mechanism. *Photosynth. Res.* **121**, 185–192.
- Wang, Y., Stessman, D.J. and Spalding, M.H. (2015) The CO₂ concentrating mechanism and photosynthetic carbon assimilation in limiting CO₂: how *Chlamydomonas* works against the gradient. *Plant J.* **82**, 429–448.
- Whitney, S.M., Houtz, R.L. and Alonso, H. (2011) Advancing our understanding and capacity to engineer nature's CO₂-sequestering enzyme, Rubisco. *Plant Physiol.* **155**, 27–35.
- Yamano, T., Tsujikawa, T., Hatano, K., Ozawa, S., Takahashi, Y. and Fukuzawa, H. (2010) Light and low-CO₂-dependent LCIB-LCIC complex localization in the chloroplast supports the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Cell Physiol.* **51**, 1453–1468.
- Yamano, T., Asada, A., Sato, E. and Fukuzawa, H. (2014) Isolation and characterization of mutants defective in the localization of LCIB, an essential factor for the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Photosynth. Res.* **121**, 193–200.
- Yamano, T., Sato, E., Iguchi, H., Fukuda, Y. and Fukuzawa, H. (2015) Characterization of cooperative bicarbonate uptake into chloroplast stroma in the green alga *Chlamydomonas reinhardtii*. *Proc. Natl Acad. Sci. USA*, **112**, 7315–7320.
- Ynalvez, R.A., Xiao, Y., Ward, A.S., Cunnusamy, K. and Moroney, J.V. (2008) Identification and characterization of two closely related beta-carbonic anhydrases from *Chlamydomonas reinhardtii*. *Physiol. Plant.* **133**, 15–26.
- Zhang, R., Patena, W., Armbruster, U., Gang, S.S., Blum, S.R. and Jonikas, M.C. (2014) High-throughput genotyping of green algal mutants reveals random distribution of mutagenic insertion sites and endonucleolytic cleavage of transforming DNA. *Plant Cell*, **26**, 1398–1409.
- Zhu, X.G., Long, S.P. and Ort, D.R. (2010) Improving photosynthetic efficiency for greater yield. *Annu. Rev. Plant Biol.* **61**, 235–261.

Supporting information

Additional Supporting information may be found in the online version of this article:

Figure S1 Expression of fluorescent-tagged CCM components in *Chlamydomonas* and tobacco (from Figure 1).

Figure S2 Co-expression of Venus-fused CCM components with Mitotracker Red CMXRos in *Chlamydomonas*.

Figure S3 Expression of GFP-fused CCM components carrying native *Arabidopsis* chloroplast transit peptides in tobacco (from Figure 3).

Figure S4 Stable expression of LCIA: GFP and HLA3: GFP in *Arabidopsis* (from Figure 5a).

Figure S5 Chlorophyll content of transgenic *Arabidopsis* plants expressing LCIA or HLA3.

Figure S6 Specific leaf area (area/DW) of transgenic LCIA or HLA3 *Arabidopsis* plants.

Figure S7 DNA sequences of codon-optimized LCIA and HLA3.

Table S1 Sequences of synthetic oligonucleotides used in this study.